



IDAHO BLM

TECHNICAL BULLETIN

The Results of Using "Hobble Creek" Sagebrush on Two Fire Rehabilitations

by
James H. Klott
and
Chris Ketchum



TECHNICAL BULLETIN 91-1
March 1991

BUREAU OF LAND MANAGEMENT
IDAHO STATE OFFICE
3380 Americana Terrace
Boise, Idaho 83706

QL
84.2
.L352
no. 91-1



27313926

D 88027341

QL
84.2
.L352
no.91-1

THE RESULTS OF USING "HOBBLE CREEK" SAGEBRUSH ON TWO FIRE REHABILITATIONS

by

James H. Klott
Wildlife Biologist
Deep Creek Resource Area
138 South Main
Malad City, Idaho 83252

and

Chris Ketchum
Wildlife Biologist
Burley District Office
200 South Oakley Highway
Route 3, Box 1
Burley, Idaho 83318

BLM LIBRARY
SC-653, BLDG. 50
DENVER FEDERAL CENTER
P. O. BOX 25047
DENVER, CO 80225-0047

TECHNICAL BULLETIN 91-1
MARCH 1991

Bureau of Land Management
Idaho State Office
3380 Americana Terrace
Boise, Idaho 83706

1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025

THE RESULTS OF USING 'HOBBLE CREEK' SAGEBRUSH ON TWO FIRE REHABILITATIONS

James H. Klott, Deep Creek Resource Area Wildlife Biologist, 138 South Main, Malad City, ID 83252

and

Chris Ketchum, Burley District Wildlife Biologist, 200 South Oakley Highway, Route 3 Box 1, Burley, ID 83318


INTRODUCTION

Two fire rehabilitation efforts within the Deep Creek Resource Area have used 'Hobble Creek' sagebrush seedlings as a method to speed the recovery of wildlife habitats following wild fires. The Grandine Fire burned approximately 360 acres on August 8, 1986, part of which was critical mule deer (Odocoileus hemionus) winter range. The Badger Hole Fire burned nearly 3,000 acres during August 9 and 10, 1988. Wildlife values impaired were sage grouse (Centrocercus urophasianus) and sharp-tailed grouse (Tympanuchus phasianellus) nesting and brood rearing habitat and mule deer transition and summer range.

'Hobble Creek' sagebrush is an accession of mountain big sagebrush (Artemisia tridentata ssp vaseyana) (Welch et al. 1986). 'Hobble Creek' sagebrush is preferred by wintering mule deer (Welch et al. 1986), and readily consumed by pygmy rabbits (Brachylagus idahoensis) (White et al. 1982), sage grouse (Welch pers comm. 1990), and domestic sheep (Ovis aries) (Welch et al. 1981 and Welch et al. 1987). 'Hobble Creek' sagebrush contains 11% by dry matter of crude protein during the winter (Welch and McArthur 1979a), which exceeds the winter maintenance requirement for mule deer (Welch and McArthur 1979b, Welch et al. 1986). Welch et al. (1986) provide a detailed description of the soils, elevation, precipitation, and other characteristics of sites where 'Hobble Creek' sagebrush is grown. Jacobson and Welch (1987) describe the methods for planting 'Hobble Creek' sagebrush seed and growing and transplanting containerized seedlings (Welch et al. 1986). This technical note presents the monitoring results on the 'Hobble Creek' sagebrush seedlings at two wild fire rehabilitations. We acknowledge Bruce L. Welch of the Intermountain research Station Shrub Sciences Laboratory for reviewing this manuscript.

STUDY AREA

Precipitation at both Grandine and Badger Hole averages around 12 to 14 inches per year. The majority of the precipitation occurs in the winter and spring as snow or rain. Summer precipitation is sporadic. Elevation at the 'Hobble Creek' planting sites were 5070 feet and 5680 feet at Grandin and Badger Hole, respectively. Soils at the site of the Grandine Fire consist of the Palisade and Hiko Peak Association. Palisade soils are loams, whereas Hiko Peak soils are gravelly loams. Slopes at



the Grandine Fire varied from 10 to <50%, with a generally western aspect. At the enclosures the slope was about 10%. Soils at the Badger Hole Fire are in the Hondo-Ridgecrest-Hades Association. Hondo soils are gravelly silt loams, Ridgecrests are very gravelly loams, whereas Hades soils are silt loams. Slopes and aspects at the Badger Hole Fire were quite variable. However, within the enclosure the aspect was north eastern and the slope varied from 5 to 15%. Grazing by livestock (cattle) was not allowed for two years following the rehabilitation of both fires. In 1989 the Grandine Pasture was grazed by cattle in the fall, but this pasture was rested in 1990. Grandine pasture is grazed in a 3 treatment rest rotation system, grazed in the spring and fall with no grazing during the summer. The Badger Hole area is grazed season long (mid-June to mid-September).


METHODS

The area planted to 'Hobble Creek' sagebrush seedlings was divided into three treatments. First, a control area was designated and was unprotected from either deer or livestock use. A second treatment consisted of a 4 strand barbed wire fence, the top strand at 38 inches, to control livestock grazing, while allowing access to wintering mule deer. The third treatment was a deer proof enclosure, having a 6 inch square mesh fence approximately 6 feet tall, topped with one strand of barbed wire at a height of 7 feet. 'Hobble Creek' sagebrush seedlings were obtained from the Shrub Sciences Laboratory in Provo, Utah part of the U.S. Forest Service Intermountain Research Station. The seedlings were one year old containerized stock that had been grown throughout the winter. Seedlings were planted in April of 1987, by BLM Burley District personnel using methods outlined in Welch et al. (1986).

All of the 'Hobble Creek' sagebrush seedlings were placed into a deer proof enclosure at the Badger Hole Fire. The enclosure fence was 7 foot high composed of 6 inch square mesh. One year old containerized seedlings were obtained from the Shrub Sciences Laboratory. These seedlings were planted by volunteers as part of a Boy Scout eagle project in late April 1989. Two other areas were hand seeded to 'Hobble Creek' sagebrush at Badger Hole on about 4 to 8 inches of snow in the fall of 1988.

Rows in each of the treatments were randomly selected, then all shrubs within a row was measured for the following characteristics. Shrub height was measured using a meter stick from the ground near the stem to the tip of the longest branch. Shrub canopy was measured with a meter stick in two directions. First at the widest and then the narrowest dimensions of the canopy through the center of the shrub. The two measurements were then averaged. Flower stalk length was measured from the tip of the flower stalk to the lowest floret on the stem on the longest flower stalk on each shrub. The number of flowering stems were counted for each shrub sampled. The extensive utilization method was used to evaluate browse use on the sampled shrubs (BLM Manual 4430.C).

Seedling survival was determined using one of two methods. At Grandine, spacing, the presence of a depressions, the presence of dead stems, and the presence of live shrubs were used to determine survival rates. However, at Badger Hole 5 permanent 30 foot radius circular plots were established and the number of sagebrush seedlings within each plot



were recorded. Additionally, mortality due to ants (*Formica* sp.) was evaluated by using 16 foot radius circular plots and measuring the distance from the plot center to each alive or dead seedling at 20 randomly selected ant mounds and 20 plots without ant mounds. Areas hand seeded to 'Hobble Creek' sagebrush were evaluated for seedling establishment with 1/100 acre circular plots.

RESULTS

Survival rates of the 'Hobble Creek' sagebrush seedlings were 88.28% in 1989 and 86.21% in 1990 at the Badger Hole enclosure. While conducting the survivals in 1989, it was noted that the majority of the mortality (65%) occurred within 15 feet of ant mounds in the permanent plots. The number of shrubs with flower stalks increased from 8.59% in 1989 to 44.80% in 1990. Based on the ant plots compared to the control plots it appears that the major cause of mortality at the Badger Hole enclosure was due to defoliation by ants (Table 1).

Hand seeding on snow resulted in an average of 49 seedlings for the plots in 1989. This increased to 69.5 seedlings per plot in 1990.

After four years, the 'Hobble Creek' sagebrush survival study at Grandine indicated that there was a significantly lower survival rate (48.88%) in the area not protected from browsing (control treatment) compared to the livestock and deer enclosures, 79.09% and 78.38%, respectively. Not only was survival impacted by browsing, but shrub height, number of flowering stems per shrub, and shrub canopy size were also influenced (Table 2). Browse utilization showed that cattle consumed nearly 35% of the current annual growth on the 'Hobble Creek' sagebrush in the control area compared to background levels of use in the two enclosures (Table 3). Wintering mule deer increased the level of use on the 'Hobble Creek' sagebrush by an additional 22% in the control and 45% within the livestock enclosure.

DISCUSSION

Data from the hand seeded areas indicates that seeding on 4 to 8 inches of snow is a successful way to establish 'Hobble Creek' sagebrush. Differences between the numbers of seedlings in 1989 and 1990 suggest that due to the small size of some sagebrush seedlings they were not detected or some of the 'Hobble Creek' sagebrush seed remained viable until the second year. Due to the location of the hand seeded areas with respect to unburned sagebrush areas, it is unlikely that the seed came from a local source. The data also suggest that aerial seeding of sagebrush in the late fall is suitable for sites where planting seed to a 5 mm depth (Jacobson and Welch 1987) is not practical.

Based on the utilization data in the fall of 1989 for the control treatment (no enclosure), it appears that cattle forage on 'Hobble Creek' sagebrush. 'Hobble Creek' sagebrush is readily consumed by other forms of livestock, particularly sheep (Welch et al 1986b). Clary (1986) noted both sheep and cattle foraged on black sagebrush (*A. nova*). Richardson et al. (1986) reported cattle readily consumed mountain big sagebrush (accession unknown) seedlings. It is unknown from this study if the season of livestock use, trampling by livestock, the amount of current

annual growth removed during the growing season, or if livestock use coupled with winter browsing by mule deer was the source of the higher mortality in the control treatment. However, research by Wright (1970), Cook and Child (1971) and Clary (1986) indicates that season of use influences mortality in big sagebrush and black sagebrush.

In addition to higher mortality rates, the different levels or season of use appeared to impact the vigor of 'Hobble Creek' sagebrush, particularly canopy diameter and reproductive potential. Hsiao (1986) and Rodriguez and Welch (1989) found similar results were caused by severe defoliation and excessive browsing, respectively. Hsiao reported that sagebrush defoliation caused higher mortality and described the symptoms of reduced vigor as shrubs producing less foliage, a smaller percentage of live branches, and fewer branches producing flower stalks. Results at Grandine, shorter flower stalk length and fewer flowering stems, were similar to those reported by Rodriguez and Welch (1989) and Wagstaff and Welch (1991) on a winter range in Utah. Mortality within the livestock and deer exclosures at Grandine appeared to be principally due to pocket gophers.

At the Badger Hole area, the majority of the mortality appeared to be due to defoliation by ants rather than pocket gophers. Seedlings planted more than 10 feet from ant mounds had a first year mortality rate (5.03%) comparable to plots without ant mounds 4.52%. A first year mortality rate (11.72%) at Badger Hole is slightly higher than other first year mortality rates (Welch pers. comm. 1990). In cases where the planting of seedlings are contracted, a stipulation should be made to plant shrub seedlings at least 10 feet from ant mounds.

Hunter et al. (1980) and Richardson et al. (1986) have recommended that areas planted to shrub seedlings should be protected from livestock grazing for several years. The results at Grandine support their conclusions, but suggest that the type of fencing should reflect the objectives of the plantings. If the objective of the plantation is to produce a seed source to speed revegetation or seed to be scattered in other areas, then a deer fences around plantations should be considered to maximize the amount of seed produced. If the objective is to provide forage for wintering mule deer, then some type of fencing should be considered to control livestock use.

Regardless of the type of fencing around a plantation, 'Hobble Creek' sagebrush seedling establishment in adjacent areas may be slow. McArthur et al. (1988) found that seedling establishment and survival was greatly reduced by excessive browsing mule deer on a winter range in Utah. They concluded that seedlings could not withstand heavy utilization. Rodriguez and Welch (1989) believed that excessive browsing reduced seed production, and therefore, reduced the potential for new seedlings.




Table 1. A comparison of plots centered in ant mounds to plots without ant mounds.

Characteristics	Ant Mounds	Control
n	20	20
X Distance to seedling (feet)	10.72	10.59
X Distance to live seedling (feet)*	12.58	10.54
X Distance to dead seedling (feet)*	6.83	11.48
No. Dead Seedlings*	43	7
No. Live Seedlings*	96	148
Survival rate (%)*	69.06	95.48

* Denotes a statistically significant difference $P < 0.05$




Table 2. Comparison of vegetative characteristics of 'Hobble Creek' sagebrush seedlings in control, livestock exclosure, and deer exclosure treatments.

Characteristic	Control (No Exclosure)	Livestock Exclosure	Deer Exclosure
Shrub height (cm)			
1989*	13.87	25.83	26.27
1990*	23.40	30.70	37.30
Canopy diameter (cm)			
1989*	12.53	23.27	27.38
1990*	21.85	31.06	39.03
Flower stalk length (cm)			
1989*	10.00	23.89	30.33
1990	17.33	17.62	22.42
No. Flower stalks			
1989*	0.30	3.53	14.87
1990*	7.36	20.47	50.57
Flowering Shrubs 1990 (%)*	55.17	83.91	96.5
Survival rate 1990 (%)*	48.88	79.09	78.38

* Denotes a statistically significant difference $P < 0.05$



Table 3. Utilization of 'Hobble Creek' sagebrush by livestock and deer in the Grandine Pasture.


Utilization	Control (No enclosure)	Livestock Enclosure	Deer Enclosure
Fall 1989*	41.66	6.00	4.67
Spring 1990*	64.33	51.67	10.33
Fall 1990	5.28	4.71	5.63

* Denotes a statistically significant difference $P < 0.05$



LITERATURE CITED

- Clary, W.P. 1986. Black sagebrush response to grazing in the east-central Great Basin. pp. 181-185. In E.D. McArthur and B.L. Welch (compilers) Proceedings--symposium on the biology of Artemisia and Chrysothamnus, 1984. Gen. Tech. Rep. INT-200. U.S. Dep. Agric. For. Serv. Intermtn. Res. Sta., Ogden, UT. 398 p.
- Cook, C.W. and R.D. Child. 1971. Recovery of desert plants in various states of vigor. *J. Range Manage.* 24:339-343.
- Hunter, R.B., A. Wallace, E.M. Wallace. 1980. Fencing enhances shrub survival and growth for Mojave desert Nevada test site. *Great Basin Nat. Memoirs.* 4:212-215.
- Hsiao, T.H. 1986. Biology and demography of the sagebrush defoliator and its impacts on big sagebrush. pp. 191-198. In E.D. McArthur and B.L. Welch (compilers) Proceedings--symposium on the biology of Artemisia and Chrysothamnus, 1984. Gen. Tech. Rep. INT-200. U.S. Dep. Agric. For. Serv. Intermtn. Res. Sta., Ogden, UT. 398 p.
- Jacobson, T.L.C. and B.L. Welch. 1987. Planting depth of 'Hobble Creek' mountain big sagebrush seed. *Great Basin Nat.* 47:497-499.
- McArthur, E.D., A.C. Blauer, and S.C. Sanderson. 1988. Mule deer induced mortality of mountain big sagebrush. *J. Range Manage.* 41:114-117.
- Richardson, B.Z., S.B. Monsen, and D.M. Bowers. 1986. Interseeding selected shrubs and herbs on mine disturbances in southeastern Idaho. pp. 134-139. In E.D. McArthur and B.L. Welch (compilers) Proceedings--symposium on the biology of Artemisia and Chrysothamnus, 1984. Gen. Tech. Rep. INT-200. U.S. Dep. Agric. For. Serv. Intermtn. Res. Sta., Ogden, UT. 398 p.
- Rodriguez, R.L. and B.L. Welch. 1989. Effects of heavy grazing by mule deer on 'Hobble Creek' mountain big sagebrush seed stalk production. pp. 141-143. In A. Wallace, E.D. McArthur, and M.R. Haferkamp. (compilers) Proceedings--symposium on shrub ecophysiology and biotechnology 1987. Gen. Tech. Rep. INT-256. U.S. Dep. Agric. For. Serv. Intermtn. Res. Sta., Ogden, UT. 183 p.
- U.S. Dep. Interior, Bur. Land Manage. Manual 4430.C Extensive Utilization Method
- Wagstaff, F.J. and B.L. Welch. 1991. Seedstalk production of mountain big sagebrush enhanced through short-term protection from heavy browsing. *J. Range Manage.* 44:72-74.
- Welch, B.L. 1990 pers. comm. Principal plant physiologist, Intermtn. Res. Sta. Shrub Science Laboratory, Provo, UT.



- Welch, B.L. and E.D. McArthur. 1979a. Variation in winter levels of crude protein among Artemisia tridentata subspecies grown in a uniform garden. J. Range Manage. 32:467-469.
- Welch, B.L. and E.D. McArthur. 1979b. Feasibility of improving big sagebrush (Artemisia tridentata) for use on mule deer winter ranges. pp. 451-473. In J.R. Goodin and D.K. Northington. eds. Proceedings arid lands international conferences on plant resources 1978. Texas Tech. Univ., Lubbock.
- Welch, B.L., E.D. McArthur, and J.N. Davis. 1981. Differential preference of wintering mule deer for accessions of big sagebrush and for black sagebrush. J. Range Manage. 34:409-411.
- Welch, B.L., E.D. McArthur, D.L. Nelson, J.C. Pederson, and J.N. Davis. 1986. 'Hobble Creek'- a superior selection of low-elevation mountain big sagebrush. U.S. Dep. Agric. For. Serv., Res. Pap. INT-370, Intermtn. Res. Sta., Ogden, Utah. 10 pp.
- Welch, B.L., E.D. McArthur, and R.L. Rodriguez. 1987. Variation in utilization of big sagbrush accession by wintering sheep. J. Range Manage. 40:113-115.
- White, S.M., J.T. Flinders, and B.L. Welch. 1982. Preference of pygmy rabbits (Brachylagus idahoensis) for various populations of big sagebrush (Artemisia tridentata). J. Range Manage. 41:114-117.
- Wright, H.A. 1970. Response of big sagebrush and three-tip sagebrush to season of clipping. J. Range Manage. 23:20-22.




Figure 1. "Hobble Creek" sagebrush seedlings (Fall, 1989). All seedlings are the same age. Note the differences in shrub size, seedstalk number, and seedstalk length. Cover board squares are 5 cm. x 5 cm.




Figure 1a. Seedling in the control treatment: no protection from deer or livestock.






Figure 1b. Seedling in the livestock enclosure.







Figure 1c. Seedling in the deer enclosure.



027341

ER'S CARD

no. 81-1

ing "Hobble
rush on two

	OFFICE	DATE RETURNED

(Continued on reverse)

BLM LIBRARY
SC-653, BLDG. 50
DENVER FEDERAL CENTER
P. O. BOX 25047
DENVER, CO 80225-0047



BLM-ID-PT- 91-009-4350